

PARTIAL DYNAMICAL SYMMETRY AND THE PHONON STRUCTURE OF CADMIUM ISOTOPES*

A. LEVIATAN, N. GAVRIELOV

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

(Received November 16, 2018)

The phonon structure and spectral properties of states in ^{110}Cd are addressed by including proton excitations in the phonon basis and exploiting a partial dynamical symmetry that mixes only certain classes of states and maintains the vibrational character in the majority of normal states.

DOI:10.5506/APhysPolB.50.561

The cadmium isotopes since long have been considered as archetypal examples of spherical vibrators, manifesting the U(5) dynamical symmetry (DS) [1]. Recent studies, however, have cast doubt on the validity of this description [2, 3]. In the present contribution, we address this question from a symmetry-oriented perspective, focusing on ^{110}Cd [4].

The U(5)-DS limit of the interacting boson model (IBM) [1], corresponds to the chain of nested algebras: $U(6) \supset U(5) \supset SO(5) \supset SO(3)$. The basis states $[[N], n_d, \tau, n_\Delta, L\rangle$ have quantum numbers which are the labels of irreducible representations of the algebras in the chain. Here, N is the total number of monopole (s) and quadrupole (d) bosons, n_d and τ are the d -boson number and seniority, respectively, and L is the angular momentum. The multiplicity label n_Δ counts the maximum number of d -boson triplets coupled to $L = 0$. The U(5)-DS Hamiltonian has the form of [1]

$$\hat{H}_{\text{DS}} = t_1 \hat{n}_d + t_2 \hat{n}_d^2 + t_3 \hat{C}_{\text{SO}(5)} + t_4 \hat{C}_{\text{SO}(3)}, \quad (1)$$

where \hat{C}_G is a Casimir operator of G , and $\hat{n}_d = \sum_m d_m^\dagger d_m = \hat{C}_{U(5)}$. \hat{H}_{DS} is completely solvable for *any* choice of parameters t_i , with eigenstates $[[N], n_d, \tau, n_\Delta, L\rangle$ and energies $E_{\text{DS}} = t_1 n_d + t_2 n_d^2 + t_3 \tau(\tau + 3) + t_4 L(L + 1)$. A typical U(5)-DS spectrum exhibits n_d -multiplets of a spherical vibrator, with enhanced connecting $(n_d + 1 \rightarrow n_d)$ E2 transitions.

* Presented at the Zakopane Conference on Nuclear Physics “Extremes of the Nuclear Landscape”, Zakopane, Poland, August 26–September 2, 2018.

The empirical spectrum of ^{110}Cd , shown in Fig. 1 (a), consists of both normal and intruder levels, the latter based on $2p-4h$ proton excitations across the $Z = 50$ closed shell. Experimentally known E2 rates are listed in Tables I, II. A comparison of the calculated spectrum [Fig. 1 (b)] and $B(\text{E}2)$ values [Table I], obtained from \hat{H}_{DS} (1), demonstrates that most normal states have good spherical-vibrator properties, and conform well with the properties of U(5)-DS. However, the measured rates for E2 decays from the non-yrast states, 0_3^+ ($n_d = 2$) and $[0_4^+, 2_5^+]$ ($n_d = 3$), reveal marked deviations from this behavior. In particular, $B(\text{E}2; 0_3^+ \rightarrow 2_1^+) < 7.9$, $B(\text{E}2; 2_5^+ \rightarrow 4_1^+) < 5$, $B(\text{E}2; 2_5^+ \rightarrow 2_2^+) = 0.7^{+0.5}_{-0.6}$ W.u. are extremely small compared to the U(5)-DS values: 46.29, 19.84, 11.02 W.u., respectively. Absolute $B(\text{E}2)$ values for transitions from the 0_4^+ state are not known, but its branching ratio to 2_2^+ is small.




Fig. 1. (a) Experimental spectrum and representative E2 rates [3, 5] (in W.u.) of normal and intruder levels (0_2^+ , 2_3^+ , 4_3^+ , 2_4^+) in ^{110}Cd . (b) Calculated U(5)-DS spectrum obtained from \hat{H}_{DS} (1) with parameters $t_1 = 715.75$, $t_2 = -t_3 = 42.10$, $t_4 = 11.38$ keV and $N = 7$. (c) Calculated U(5)-PDS-CM spectrum, obtained from \hat{H} (3) with parameters $t_1 = 767.83$, $t_2 = -t_3 = 73.62$, $t_4 = 18.47$, $r_0 = 2.15$, $e_0 = -6.92$, $\kappa = -72.73$, $\Delta = 9978.86$, $\alpha = -42.78$ keV and $N = 7$ (9) in the normal (intruder) sector. (d) Classes of low-lying U(5)-DS states.

TABLE I

Absolute (relative in square brackets) $B(E2)$ values in W.u. for E2 transitions from normal levels in ^{110}Cd . The experimental (EXP) values are taken from [3, 5]. The U(5)-DS [U(5)-PDS-CM] values are obtained for an E2 operator $e_B \hat{Q} [e_B^{(N)} \hat{Q}^{(N)} + e_B^{(N+2)} \hat{Q}^{(N+2)}]$ with $e_B = 1.964$ [$e_B^{(N)} = 1.956$ and $e_B^{(N+2)} = 1.195$] (W.u.) $^{1/2}$, where $\hat{Q} = d^\dagger s + s^\dagger \tilde{d}$ and $\hat{Q}^{(N)}$ denotes its projection onto the $[N]$ boson space. In both calculations, the boson effective charges were fixed by the empirical $2_1^+ \rightarrow 0_1^+$ rate. Intruder states $0_{2;i}^+$, $2_{3;i}^+$, $4_{3;i}^+$, $2_{4;i}^+$, are marked by a subscript 'i'. ^aFrom Ref. [3].

L_i	L_f	EXP	U(5)-DS	U(5)-PDS-CM
2_1^+	0_1^+	27.0 (8)	27.00	27.00
4_1^+	2_1^+	42 (9)	46.29	45.93
2_2^+	2_1^+	30 (5); 19 (4) ^a	46.29	46.32
	0_1^+	1.35 (20); 0.68 (14) ^a	0.00	0.00
0_3^+	2_2^+	< 1680 ^a	0.00	55.95
	2_1^+	< 7.9 ^a	46.29	0.25
6_1^+	4_1^+	40 (30); 62 (18) ^a	57.86	55.30
	4_2^+	< 5 ^a	0.00	0.00
	$4_{3;i}^+$	14 (10); 36 (11) ^a		2.39
4_2^+	4_1^+	12_{-6}^{+4} ; $^{a}10.7_{-4.8}^{+4.9}$	27.55	27.45
	2_2^+	32_{-14}^{+10} ; 22 (10) ^a	30.31	30.03
	2_1^+	$0.20_{-0.09}^{+0.06}$; 0.14 (6) ^a	0.00	0.00
	$2_{3;i}^+$	< 0.5 ^a		0.005
3_1^+	4_1^+	$5.9_{-4.6}^{+1.8}$; $^{a}2.4_{-0.8}^{+0.9}$	16.53	16.48
	2_2^+	32_{-24}^{+8} ; 22.7 (69) ^a	41.33	41.12
	2_1^+	$1.1_{-0.8}^{+0.3}$; 0.85 (25) ^a	0.00	0.00
	$2_{3;i}^+$	< 5 ^a		0.012
0_4^+	2_2^+	[< 0.65 ^a]	57.86	1.24
	2_1^+	[0.010 ^a]	0.00	31.76
	$2_{3;i}^+$	[100 ^a]		16.32
2_5^+	0_3^+	24.2 (22) ^a	27.00	22.28
	4_1^+	< 5 ^a	19.84	0.19
	2_2^+	$^{a}0.7_{-0.6}^{+0.5}$	11.02	0.12
	2_1^+	$2.8_{-1.0}^{+0.6}$	0.00	0.00
	$2_{3;i}^+$	< 5 ^a		0.002
	$0_{2;i}^+$	< 1.9 ^a		0.20

Attempts to explain the above deviations in terms of mixing between the normal spherical [U(5)-like] states and intruder deformed [SO(6)-like] states have been shown to be unsatisfactory [2, 3]. This has led to the conclusion that the normal-intruder strong-mixing scenario needs to be rejected, and

TABLE II

$B(E2)$ values (in W.u.) for E2 transitions from intruder levels in ^{110}Cd . Notation and relevant information on the observables shown are as in Table I.

L_i	L_f	EXP	U(5)-PDS-CM
$0_{2;i}^+$	2_1^+	$< 40^a$	14.18
$2_{3;i}^+$	$0_{2;i}^+$	$29 (5)^a$	29.00
	0_1^+	$0.31_{-0.12}^{+0.08}; 0.28 (4)^a$	0.08
	2_1^+	$0.7_{-0.4}^{+0.3}; {}^a0.32_{-0.14}^{+0.10}$	0.00
	2_2^+	$< 8^a$	0.96
$2_{4;i}^+$	2_1^+	$0.019_{-0.019}^{+0.020}$	0.10
$4_{3;i}^+$	2_1^+	$0.22_{-0.19}^{+0.09}; 0.14 (4)^a$	0.49
	2_2^+	$2.2_{-2.2}^{+1.4}; 1.2(4)^a$	0.00
	$2_{3;i}^+$	$120_{-110}^{+50}; 115 (35)^a$	42.62
	4_1^+	$2.6_{-2.6}^{+1.6}; {}^a1.8_{-1.5}^{+1.0}$	0.00

have raised serious questions on the appropriateness of the multi-phonon interpretation [2, 3]. In what follows, we consider a possible explanation for the ‘‘Cd problem’’, based on U(5) partial dynamical symmetry (PDS). The latter corresponds to a situation in which the U(5)-DS is obeyed by only a subset of states and is broken in other states [6]. Similar PDS-based approaches have been implemented in nuclear spectroscopy, in conjunction with the SU(3)-DS [7–9] and SO(6)-DS [10, 11] chains of the IBM.

As depicted in Fig. 1 (d), the lowest spherical-vibrator levels comprise three classes of states. Specifically, Class A: $n_d = \tau = 0, 1, 2, 3$ ($n_\Delta = 0$); Class B: $n_d = \tau + 2 = 2, 3$ ($n_\Delta = 0$); Class C: $n_d = \tau = 3$ ($n_\Delta = 1$). In the U(5)-DS calculation of Fig. 1 (b), applicable to normal states only, the ‘‘problematic’’ states [0_3^+ ($n_d = 2$) and 2_5^+ ($n_d = 3$)] belong to class B, and 0_4^+ ($n_d = 3$) belongs to class C. The remaining ‘‘good’’ spherical-vibrator states [0_1^+ ($n_d = 0$); 2_1^+ ($n_d = 1$); $4_1^+, 2_2^+$ ($n_d = 2$); $6_1^+, 4_2^+, 3_1^+$ ($n_d = 3$)] belong to class A. As mentioned, the spherical-vibrator interpretation is valid for most normal states in Fig. 1 (a), but not all. We are thus confronted with a situation in which some states in the spectrum (assigned to class A) obey the predictions of U(5)-DS, while other states (assigned to classes B and C) do not. These empirical findings signal the presence of U(5)-PDS.

The construction of a Hamiltonian with U(5)-PDS follows the general algorithm [6] and leads to the form of

$$\hat{H}_{\text{PDS}} = \hat{H}_{\text{DS}} + r_0 G_0^\dagger G_0 + e_0 \left(G_0^\dagger K_0 + K_0^\dagger G_0 \right), \quad (2)$$

where $G_0^\dagger = [(d^\dagger d^\dagger)^{(2)} d^\dagger]^{(0)}$, $K_0^\dagger = s^\dagger (d^\dagger d^\dagger)^{(0)}$. The last two terms in Eq. (2) annihilate the states $||[N], n_d = \tau, \tau, n_\Delta = 0, L\rangle$ with $L = \tau, \tau + 1, \dots, 2\tau - 2, 2\tau$.

These states, which include those of class A, form a subset of U(5) basis states, hence remain solvable eigenstates of \hat{H}_{PDS} (2) with good U(5) symmetry. It should be noted that while \hat{H}_{DS} (1) is diagonal in the U(5)-DS chain, the r_0 and e_0 terms can connect states with different n_d and/or τ . Accordingly, the remaining eigenstates of \hat{H}_{PDS} (2), in particular those of classes B and C, are mixed with respect to U(5) and SO(5). The U(5)-DS is thus preserved in a subset of eigenstates, for any choice of parameters in \hat{H}_{PDS} , but is broken in others. By definition, \hat{H}_{PDS} exhibits U(5)-PDS.

The combined effect of normal and intruder states can be studied within the interacting boson model with configuration mixing (IBM-CM) [12]. The Hamiltonian for the two configurations has the form of [4]

$$\hat{H} = \hat{H}_{\text{PDS}}^{(N)} + \hat{H}_{\text{intrud}}^{(N+2)} + \hat{V}_{\text{mix}}. \tag{3}$$

For ^{110}Cd , the Hamiltonian in the normal sector is taken to be \hat{H}_{PDS} of Eq. (2), projected onto a space of $N = 7$ bosons. The SO(6)-type of Hamiltonian in the intruder sector is $\hat{H}_{\text{intrud}} = \kappa \hat{Q} \cdot \hat{Q} + \Delta$, projected onto a space of $N = 9$ bosons. $\hat{V}_{\text{mix}} = \alpha[(s^\dagger)^2 + (d^\dagger d^\dagger)^{(0)}] + \text{H.c.}$ is a mixing term between the two spaces. In general, an eigenstate of \hat{H} , $|\Psi\rangle = a|\Psi_n^{(N)}\rangle + b|\Psi_i^{(N+2)}\rangle$, involves a mixture of normal (n) and intruder (i) components with N and $N+2$ bosons, respectively.

As seen in Fig. 1 (c) and Tables I, II, the IBM-PDS-CM calculation provides a good description of the empirical data in ^{110}Cd . The U(5) decomposition of the resulting eigenstates is shown in Fig. 2. The normal states of class A retain good U(5) symmetry to a good approximation. Their $|\Psi_n^{(N)}\rangle$ part involves a single n_d -component. The mixing with the intruder states is weak (small b^2) of the order of a few percent. The high degree of purity is reflected in the calculated $B(\text{E}2)$ values for transitions between class A states, which are very similar to those of U(5)-DS. In contrast, the structure of the non-yrast states assigned originally to classes B and C, whose decay properties show marked deviations from the U(5)-DS limit, changes dramatically. Specifically, the 0_3^+ and 0_4^+ states, which in the U(5)-DS classification are members of the two-phonon triplet and three-phonon quintuplet, interchange their character, and the U(5) decomposition of their $|\Psi_n^{(N)}\rangle$ parts peaks at $n_d = 3$ and $n_d = 2$, respectively. Similarly, for the 2_5^+ state, which is originally a member of the three-phonon quintuplet, the $|\Psi_n^{(N)}\rangle$ part now exhibits a peak at $n_d = 4$. The calculated $B(\text{E}2; 0_3^+ \rightarrow 2_1^+) = 0.25$, $B(\text{E}2; 2_5^+ \rightarrow 4_1^+) = 0.19$ and $B(\text{E}2; 2_5^+ \rightarrow 2_2^+) = 0.12$ W.u. are consistent with the measured upper limits: 7.9, 5 and $0.7_{-0.6}^{+0.5}$ W.u., respectively. The vibrational interpretation is thus maintained in the majority of low-lying normal states in ^{110}Cd .




Fig. 2. U(5) n_d -decomposition of Class A (lower panel), Classes B, C (middle panel) and intruder (upper panel) states in the U(5)-PDS-CM calculation of Fig. 1 (c). In each panel, the left(right)-hand side displays the n_d -probabilities for the normal (intruder) components of the total wave function $|\Psi\rangle = a|\Psi_n^{(N)}\rangle + b|\Psi_1^{(N+2)}\rangle$.

This work was done in collaboration with J.E. García-Ramos (Huelva) and P. Van Isacker (GANIL) and is supported by the Israel Science Foundation (Grant 586/16).

REFERENCES

- [1] F. Iachello, A. Arima, *The Interacting Boson Model*, Cambridge University Press, Cambridge 1987.
- [2] P.E. Garrett, K.L. Green, J.L. Wood, *Phys. Rev. C* **78**, 044307 (2008).
- [3] P.E. Garrett *et al.*, *Phys. Rev. C* **86**, 044304 (2012).
- [4] A. Leviatan, N. Gavriellov, J.E. García-Ramos, P. Van Isacker, *Phys. Rev. C* **98**, 031302(R) (2018).
- [5] G. Gürdal, F.G. Kondev, *Nucl. Data Sheets* **113**, 1315 (2012).
- [6] A. Leviatan, *Prog. Part. Nucl. Phys.* **66**, 93 (2011).
- [7] A. Leviatan, *Phys. Rev. Lett.* **77**, 818 (1996).
- [8] A. Leviatan, I. Sinai, *Phys. Rev. C* **60**, 061301(R) (1999).
- [9] A. Leviatan, J.E. García-Ramos, P. Van Isacker, *Phys. Rev. C* **87**, 021302(R) (2013).
- [10] J.E. García-Ramos, A. Leviatan, P. Van Isacker, *Phys. Rev. Lett.* **102**, 112502 (2009).
- [11] A. Leviatan, N. Gavriellov, *Phys. Scr.* **92**, 114005 (2017).
- [12] P.D. Duval, B.R. Barrett, *Nucl. Phys. A* **376**, 213 (1982).